

FRAME-DRAGGING, DISK WARPING, JET PRECESSING, AND DIPPED X-RAY LIGHTCURVE OF SW J1644+57

WEI-HUA LEI^{1,2}, BING ZHANG¹, HE GAO¹

Draft version February 21, 2012

ABSTRACT

The X-ray transient source Sw J1644+57 recently discovered by Swift is believed to be triggered by tidal disruption of a star by a rapidly spinning supermassive black hole (SMBH). For such events, the outer disk is very likely misaligned with respect to the equatorial plane of the spinning SMBH, since the incoming star before disruption most likely has an inclined orbital plane. The tilted disk is subject to the Lense-Thirring torque, which tends to twist and warp the disk due to the Bardeen-Petterson effect. The inner disk tends to align with the SMBH spin, while the outer region tends to remain in the stellar orbital plane, with a transition zone around the Bardeen-Petterson radius. The relativistic jet launched via the Blandford-Znajek mechanism from the spinning SMBH would undergo precession. The X-ray lightcurve of Sw J1644+57 shows a quasi-periodic (2.7-day) variation with noticeable narrow dips. We numerically solve a warping disk solution and propose a jet-processing model by invoking a Blandford-Znajek jet collimated by a wind launched near the Bardeen-Petterson radius. Through simulations, we show that the narrow dips in the X-ray lightcurve can be reproduced for a range of geometric configurations. From data we infer that the inclination angle of the initial stellar orbit is in the range of $10^\circ - 20^\circ$ from the SMBH equatorial plane, that the jet should have a moderately high Lorentz factor, and that the inclination angle, jet opening angle, and observer's viewing angle are such that the duty cycle of the line-of-sight sweeping the jet cone is somewhat less than 0.5.

Subject headings: accretion, accretion disks—black hole physics—magnetic fields

1. INTRODUCTION

The discover of the hard X-ray transient event Swift J16449.3+573451 (“Sw J1644+57” hereafter, Burrows et al. 2011) by the Swift satellite (Gehrels et al. 2004) has stimulated a great interest in studying jets launched from tidal disruption events (TDEs). The long variability time scale $\delta t \sim 100$ s (Burrows et al. 2011) and its location near the center of a $z = 0.354$ host galaxy (Levan et al. 2011) link Sw J1644+57 to a super-massive black hole (SMBH) (Burrows et al. 2011; Bloom et al. 2011). The sharp onset and gradual fade-away of X-ray flux refer to tidal disruption of a star by a dormant SMBH (Bloom et al. 2011; Burrows et al. 2011). The super-Eddington X-ray luminosity (Burrows et al. 2011), bright radio afterglow (Zauderer et al. 2011), as well as a stringent historical X-ray flux upper limit suggest that a relativistic jet is launched from a SMBH during the TDE, which is not expected in most previous TDE studies (e.g. Kobayashi et al. 2004; Strubbe & Quataert 2010; Lodato & Rossi 2011, but see Lu et al. 2008; Gao et al. 2010; Giannios & Metzger 2011). Modelling the emission of Sw J1644+57 suggests that the jet is highly “particle starved” (Burrows et al. 2011), favoring a magnetically launched jet, likely launched via the Blandford-Znajek (1977, hereafter BZ) mechanism (Lei & Zhang 2011), which extracts the spin energy of the BH through a magnetic field connecting the BH event horizon and

a remote astrophysical load. From observational constraints, Lei & Zhang (2011) found that the SMBH in Sw J1644+57 carries a moderate to high spin.

For such events, since the star initially has no knowledge about the BH spin orientation before being disrupted and accreted, it is most likely that the initial stellar orbit is mis-aligned with the equatorial plane of the spinning BH. A natural expectation is that at least the outer part of the accretion disk is also misaligned. The tilted disk surrounding a spinning BH is subject to the Lense-Thirring (Lense & Thirring 1918, LT hereafter) torque. The combined action of the LT torque and the internal viscosity of the accretion disk would lead to a twisted and warped disk, with the inner part of the disk bent towards the BH equatorial plane due to the frame-dragging effect. This is known as the Bardeen-Petterson effect (Bardeen & Petterson 1975, hereafter BP). For a fully developed BP disk, the inner part of the disk tends to be aligned with the BH equatorial plane, while the outer part of the disk remains aligned with the original stellar orbital plane. The transition radius between these two regimes is the BP radius. The warped disks have been directly observed by water maser observations in NGC 4258 (Miyoshi et al. 1995; Neufeld & Maloney 1995; Herrnstein, Greenhill & Moran 1996) and the Circinius galaxy (Greenhill et al. 2003). The apparent lack of correlation between the direction of the radio jets emanating from active galactic nuclei (AGN) and the plane of the host galaxy disks (Kinney et al. 2000; Schmitt et al. 2002) can also be explained by disk warping.

Besides this theoretical motivation, some curious observational facts also point towards the possibility of a warped disk. Burrows et al. (2011) reported a rough

¹ Department of Physics and Astronomy, University of Nevada Las Vegas, 4505 Maryland Parkway, Box 454002, Las Vegas, NV 89154-4002, USA. Email: leiwh@physics.unlv.edu; zhang@physics.unlv.edu

² School of Physics, Huazhong University of Science and Technology, Wuhan, 430074, China

period of 230 ks (~ 2.7 days) in the long-term X-ray lightcurve, although with a $< 3\sigma$ significance. The similar rough period was also revealed recently by Saxton et al. (2012). More importantly, a closer inspection suggests that the late X-ray lightcurve show interesting dips. A natural mechanism to account for these dips would be a precessing jet, possibly caused by a warped BP disk. Stone & Loeb (2012) recently also discussed the similar topic, but didn't pay attention to these peculiar observational effects. They led to the conclusion that the stellar orbital plane is almost aligned with the BH equatorial plane due to the lack of observational evidence of jet processing.

In this paper, we develop a model to interpret the quasi-periodic feature and the narrow dips observed in the lightcurve of Sw J1644+57 within the framework of the BP mechanism. In Section 2, we first apply a mathematical method, i.e. Stepwise Filter Correlation (SFC) method (Gao et al. 2012), to study the X-ray lightcurve data of Sw 1644+57. We confirm that a ~ 2.7 -day quasi-periodic signal exists in the late lightcurve after the initial flaring phase. We then develop a numerical model in Section 3 to study disk warping and jet processing. In Section 4, we apply the observational data to constrain the model parameters, and found several parameter sets that can roughly reproduce the observations. Our conclusions are presented in Section 5.

2. OBSERVATIONAL EVIDENCE OF JET PRECESSION

Even though not with high significance, two pieces of evidence hint towards the possibility of a precessing jet in Sw J1644+57. The first evidence is the noticeable dips in the XRT lightcurve at late times. The upper panel of Fig.1 shows the XRT lightcurve of Sw J1644+57 after 5 days since the first trigger. Some deep dips are clearly visible. The dashed vertical lines mark the 2.7-day period (see more below). One can see that the dips roughly track this period. Dips are not easy to explain within models invoking emission, but would be naturally interpreted within the frame work of jet precession where the line of sight is allowed to move away from the emission beam.

The second piece of evidence is the rough 2.7-day periodicity, which was pointed out by Burrows et al. (2011) and confirmed by Saxton et al. (2012). We investigate this independently using a new method, Stepwise Filter Correlation (SFC) method, recently developed in our group (Gao et al. 2012). The SFC method is a signal processing algorithm which stepwise filters signals above a frequency, and looks for correlation between the lightcurves of adjacent filterings. If there is a spectral component around a particular frequency, the lightcurves before and after filtering this particular frequency can be very different, leading to a dip in the correlation curve. The detail of this method and its application to GRB lightcurves are presented in Gao et al. (2012). Here we apply the method to the XRT lightcurve of Sw J1644+57. We first perform an analysis to the entire lightcurve (curve (a) of lower panel of Fig.1). Since early lightcurve displays many erratic, bright X-ray flares, the signal is dominated by these high-frequency components, and the 2.7-day component cannot be identified. These flares may be related to fallback of the stellar debris with high and fluctuant accretion rate (Cannizzo et al. 2011). If we remove the early data (say, the first two days), the

2.7-day dip clearly shows up (curve (b) of lower panel of Fig.1). We then gradually remove more and more early data points and everytime redo the SFC analysis (curves (c-e) of the lower panel of Fig.1). The 2.7-day dip is always there. Our analysis suggests that, consistent with the results of Burrows et al. (2011) and Saxton et al. (2012), that the late time X-ray lightcurve of Sw J1644+57 has a quasi-periodic signal with a period roughly 2.7 days.




FIG. 1.— Upper panel: Swift XRT light curve of Sw J1644+57 from 5-30 days, which show clear dips. The dashed line shows the 2.7-day period. Data is from Burrows et al. (2011). Bottom panel: the stepwise filter correlation (SFC) analysis of the X-ray lightcurve following Gao et al. (2012). The curve (a) is the result for the entire lightcurve, curve (b), (c), (d), (e) and (f) are the analyses by manually removing the first 2, 5, 10, 15, and 20 days of the data, respectively. A dip corresponding to ~ 2.7 days is evident once the first 2-day of data is removed.

3. WARPED ACCRETION DISK AND JET PRECESSION

In simple physical terms, disruption of a star occurs when it comes to a SMBH closer than the tidal disruption radius R_T , which is determined by demanding that the mean density of space volume enclosed by R_T , i.e., $M_\bullet/(4\pi R_T^3/3)$, is equal to the density of the star, where $M_\bullet$ is the mass of the BH. The tidal disruption radius is then given by

$$R_T \simeq \left(\frac{M_\bullet}{M_*}\right)^{1/3} R_* \simeq 7 \times 10^{12} m_*^{-1/3} r_* M_{\bullet,6}^{1/3} \text{ cm} \\ \simeq 47 R_g m_*^{-1/3} r_* M_{\bullet,6}^{-2/3}, \quad (1)$$




FIG. 2.— The sketch of the model. A star is tidally disrupted as it gets close to a dormant SMBH. The star orbit is generally inclined with the SMBH spin vector $\mathbf{J}_{\text{BH}}$. After disruption, an accretion disk form, and is warped due to the Bardeen-Peterson effect. The jet is expected to precess with an angle θ_i around the BH spin axis. For a jet with half opening angle θ_j and an observer located at θ_{obs} from the BH spin axis, the angle ψ between the observer's line of sight $\hat{\mathbf{r}}_{\text{obs}}$ and jet axis $\hat{\mathbf{r}}_{\text{jet}}$ is changing with time. The line of sight can exit the jet cone at certain phases, leading to dips in the lightcurve. The Bardeen-Peterson radius R_{BP} is marked. The jet direction is defined by the disk normal direction around R_{BP} .

where $m_* = M_*/M_\odot$, $r_* = R_*/R_\odot$, $M_{\bullet,6} = M_\bullet/10^6 M_\odot$ and $R_g = GM_\bullet/c^2$. For a solar-like star ($M_* = M_\odot$ and $R_* = R_\odot$) disrupted by a $2 \times 10^6 M_\odot$ BH, the tidal disruption radius is approximately $30R_g$ from the BH.

Stellar disruption occurs if $R_{\text{ms}} \leq R_P \leq R_T$, where R_P is the periastron distance from the star, and R_{ms} is the marginally stable orbit (Bardeen et al. 1972). The star should be swallowed whole instead of being disrupted if it goes inside R_{ms} . We define $\kappa \equiv R_P/R_T$.

We idealize the disk as extending from R_{ms} to the circularization radius $R_{\text{out}} \simeq \eta R_P$, where $\eta \simeq 2$. In our calculation, the typical values of the parameters are $\eta = 2$, $m_* = 1$, $r_* = 1$, and $\kappa = 1$.

The dragging of the inertial frame (frame dragging) produced by a Kerr black hole causes precession of a particle if its orbital plane is inclined with respect to the equatorial plane of the black hole. This effect is known as Lense-Thirring precession. The precession angular velocity Ω_{LT} is given by (e.g. Wilkins 1972)

$$\Omega_{\text{LT}}(R) = \frac{2G}{c^2} \frac{J_\bullet}{R^3}, \quad (2)$$

where $J_\bullet = a_\bullet GM_\bullet^2/c$ is the BH angular momentum, and $a_\bullet$ is the BH spin parameter.

The precession period τ_p can be estimated as

$$\tau_p = 2\pi/\Omega_{\text{LT}} \quad (3)$$

For a BH with mass $2 \times 10^6 M_\odot$ and spin $a_\bullet = 0.9$, we have $\tau_p \simeq 2.7$ days at $R = 19R_g$.

When an accretion disk does not align with the equatorial plane of the BH, i.e., when the angular momentum of the accretion disk $\mathbf{L}$ is misaligned with respect to the direction of $\mathbf{J}_\bullet$, the LT effect causes a precession effect that twists the disk plane due to the coupling of $\mathbf{J}_\bullet$ with $\mathbf{L}$.

The torque tends to align the angular momentum of the matter in the disk with that of the BH, thus causing the inclination angle between the angular momentum vectors to decrease with decreasing distance from the BH. This is the BP effect (Bardeen & Petterson 1975), which is the combination of the LT effect and the internal viscosity of the accretion disk. The effect is more prominent in the inner part of the disk due to the short range of the LT effect. The outer part of the disk tends to remain in the orientation defined by the original stellar orbit.

Considering that the disk may be thick, Stone & Loeb (2012) suggested that the disk is not subject to the BP warping effect. This conclusion is based on the GRMHD simulations by Fragile et al. (2007), which invoked a weak magnetic field to account for angular momentum transport. The dynamical viscosity and magnetic stress in the disk are subject to large uncertainty, which can be larger than invoked by Fragile et al. (2007). For relativistic twisted disk around a Kerr BH, Zhuravlev & Ivanov (2008) found that BP effect can take place if the viscosity parameter α is sufficiently large. Motivated by the observational evidence discussed in Section 2, in this paper we suggest that the BP effect indeed plays a role, at least after the initial flares when the accretion rate drops significantly. To produce the high luminosity at early phase, the accretion rate is around $10^{-6} M_\odot s^{-1}$ (Lei & Zhang 2011; Shao et al. 2011). After this phase, the accretion rate declines to $\sim 10^{-8} M_\odot s^{-1}$, about 2 orders lower. For a BH with mass $10^7 M_\odot$ and the disk efficiency about 10%, the Eddington accretion rate is around $\sim 10^{-8} M_\odot s^{-1}$. Therefore we could assume that the disk becomes thin after the flare phase, and the BP effect would come into play to warp the disk.

For a warped disk, we follow the formalism of Pringle (1992). There are two viscosity parameters ν_1 and ν_2 , where ν_1 is the standard shear viscosity in a flat disk, and ν_2 is the viscosity associated with the vertical shear motion describing the diffusion of warping distortion through the disk. In order to study the BP effect, We rewrite Pringle's equation by adding an effective coupling due to the LT precession, $\Omega_{\text{LT}} \times \mathbf{L}$ (Scheuer & Feiler 1996; Lodato & Pringle 2006):

$$\begin{aligned} \frac{\partial \mathbf{L}}{\partial t} = & \frac{3}{R} \frac{\partial}{\partial R} \left[\frac{R^{1/2}}{\Sigma} \frac{\partial}{\partial R} (\nu_1 \Sigma R^{1/2}) \mathbf{L} \right] \\ & + \frac{1}{R} \frac{\partial}{\partial R} [(\nu_2 R^2) \frac{\partial \mathbf{L}}{\partial R}] - \frac{3}{2} \nu_1 \mathbf{L} \\ & + \frac{1}{R} \frac{\partial}{\partial R} \left(\frac{1}{2} \nu_2 R L \frac{\partial \mathbf{L}}{\partial R} \right) + \Omega_{\text{LT}} \times \mathbf{L}, \end{aligned} \quad (4)$$

where $L = |\mathbf{L}| = \Sigma \sqrt{GM_\bullet R}$ is the angular momentum per unit area of disk, Σ is the surface density, $\mathbf{l}$ is the unit vector indicating the local direction of the specific angular momentum in the disk. Note that R is a spherical coordinate, and is not the cylindrical radius.

The transition radius between the two regimes is known as the Bardeen-Peterson radius R_{BP} , which is the radius at which the warping propagation time-scale $t_{\nu_2} = R^2/\nu_2$ equals the local forced precession rate τ_p , giving (Scheuer & Feiler 1996; Caproni et al. 2007)

$$R_{\text{BP}} = \sqrt{\frac{\nu_2}{\Omega_{\text{LT}}(R_{\text{BP}})}}. \quad (5)$$

Here again ν_2 is the viscosity normal to the accretion disk. Roughly speaking, the disk becomes aligned with the BH spin at $R \ll R_{\text{BP}}$ and keeps its original inclination for $R \gg R_{\text{BP}}$.

Scheuer & Feiler (1996) have found an analytical steady-state solution to the above equation (removing the time-dependent term $\partial \mathbf{L} / \partial t$), subject to various simplifying assumptions about the disk density distribution and the inner boundary condition. This solution may be inaccurate at very small radii from the BH. It is based on a first-order approximation and a small inclination angle ($\theta_{\text{orbit}} \ll 1$) assumption. In our modeling, instead of applying this solution, we solve the equations numerically following the numerical scheme outlined in Pringle (1992).

The relation between ν_1 and ν_2 is the main uncertainty of the problem. For simplicity, we take $\nu_1 / \nu_2 = 1$ in the calculation. The solution for the inclination angle at time t and radius R depends on the BP radius R_{BP} , the outer disk boundary R_{out} and the orbital inclination θ_{orbit} . In order to set up a twisted disk, we add matter with an inclination angle θ_{orbit} to the grid near the outer boundary R_{out} . We run the code for about 10^5 time-steps without the LT effect to settle the disk into a numerically equilibrium state. We then evolve the disk by applying the forced precession due to the LT torque. The disk twisting evolution slows down rapidly. After about 2×10^6 runs, the inclination no longer evolve significantly, and we take this as the steady state. In Fig.3, the numerical solutions in the steady state are shown for different R_{BP} , R_{out} and θ_{orbit} .




FIG. 3.— Disk inclination θ_{inc} vs. radius R for different R_{BP} , R_{out} and θ_{orbit} : Thick solid line: $R_{\text{BP}} = 10R_g$, $R_{\text{out}} = 50R_g$ and $\theta_{\text{orbit}} = 30^\circ$; thin solid line: $R_{\text{BP}} = 10R_g$, $R_{\text{out}} = 60R_g$ and $\theta_{\text{orbit}} = 30^\circ$; dashed line: $R_{\text{BP}} = 25R_g$, $R_{\text{out}} = 60R_g$ and $\theta_{\text{orbit}} = 30^\circ$; and dotted line: $R_{\text{BP}} = 25R_g$, $R_{\text{out}} = 60R_g$ and $\theta_{\text{orbit}} = 20^\circ$.

From Fig.3, we find that the inclination of the inner disk critically depends on the BP radius R_{BP} and the initial inclination of stellar orbit θ_{orbit} . The effect of the outer disk boundary is not significant for the inner disk.

A precessing disk would induce jet precession. In our model, the relativistic jet is launched near the BH by the BZ process (Lei & Zhang 2011). However, the jet di-

rection is not determined by the normal direction at the inner disk, but is rather defined by normal direction at an outer region in the disk near R_{BP} . This is because the magnetic field threading the disk would drive an outflow from the disk surface via the Blandford & Payne (1982) mechanism. A centrifugally driven outflow is possible if the poloidal component of the magnetic field makes an angle less than 60° with the disk surface. This is most likely to occur at R_{BP} , where the disk inclination angle significantly changes. A similar proposal was introduced to study other BH accretion systems (e.g. Begelman et al. 2006) and gamma-ray bursts (e.g. Lei et al. 2007). In the following calculation, we suggest that the precession period and inclination angle of the relativistic jet is defined by the physical conditions at R_{BP} .

4. OBSERVATIONAL CONSTRAINTS

Some constraints on the model can be derived from the observational data outlined in Section 2.

The first constraint is the 2.7-day quasi-period. In the precession model delineated above, this period should correspond to the LT period at the BP radius, R_{BP} . According to Lei & Zhang (2011), the most probable value of the BH spin in Sw J1644+57 is $a_\bullet = 0.9$. We take this value for the BH spin in the calculation. Applying $\tau_p = 2.7$ days into Eq.(3), we can estimate the BP radius R_{BP} that satisfy the observations (Fig.4). For comparison, we also show the disk outer edge R_{out} in Fig.4, where $M_* = M_\odot$, $R_* = R_\odot$, $\kappa = 1$ and $\eta = 2$ are adopted as the typical values. It is found that R_{BP} is located close to the BH.




FIG. 4.— The constraint on the Bardeen-Petterson radius R_{BP} (solid line) as a function of BH mass, based on the requirement of $\tau_p(R_{\text{BP}}) = 2.7$ days with $a_\bullet = 0.9$. The disk outer edge R_{out} is also shown as the dashed line.

The next observational constraint comes from the observed shape and depth of the dips seen in the lightcurve. They are defined by the complicated combinations of several unknown parameters, including the bulk Lorentz factor Γ , the jet half opening angle θ_j , the inclination angle of the jet θ_i , as well as the observer's viewing angle θ_{obs} with respect to the BH spin axis. In the following we discuss these constraints in detail.

At any time t , the angle between the observer $\hat{r}_{\text{obs}}$ and jet axis $\hat{r}_{\text{jet}}(t)$ (the axis of the presumed conical jet) is denoted by $\psi(t)$ (see Fig.2), which is defined as

$$\psi(t) = \cos^{-1}(\hat{r}_{\text{obs}} \cdot \hat{r}_{\text{jet}}(t)). \quad (6)$$

A dip in the lightcurve can be seen if the line of sight is outside the jet cone, i.e., $\psi(t) > \theta_j$.

For simplicity, we start with a uniform conical jet, so that when $\psi < \theta_j$ the observer receives a uniform flux. For an off-axis observer at $\psi > \theta_j$, the observed flux density can be written as (Granot et al. 2002)

$$F_\nu(\psi, t) = D^3 F_{\nu/D}(0, Dt), \quad (7)$$

where $D = (1 - \beta)/(1 - \beta \cos \theta)$, $\beta = \sqrt{1 - 1/\Gamma^2}$ and $\psi = \max(0, \theta_{\text{obs}} - \theta_j)$. Define $F(\psi, t) = \nu F_\nu(\psi, t)$, we have the X-ray flux at ψ

$$F(\psi, t) = D^4 F(0, Dt). \quad (8)$$

For a tidal disruption event, the long-term lightcurve is defined by the fall-back accretion rate, which is $\propto t^{-5/3}$. The long term X-ray lightcurve of Sw J1644+57 is indeed consistent with this behavior, i.e. $F(t) \propto t^{-5/3}$. We therefore have

$$F(\psi, t) = D^{7/3} F(0, t). \quad (9)$$

The shape the lightcurve depends on several factors. The first one is the Lorentz factor. In Fig.5 we show how the Lorentz factor affects the observed structure of jet. Even if the physical jet has a sharp cutoff near the edge, the observed jet structure has a wing defined by the gradual decrease of the Doppler factor. If the jet Lorentz factor is large, the flux drops sharply as the line of sight goes away from the jet cone, and one would see a very deep dip. On the other hand, the dip would be very shallow if the jet has a smaller Lorentz factor. In the observed lightcurve, the dips do not have uniform depth, and in some periods the dips disappear completely. This might be caused by fluctuation of the jet Lorentz factor as a function of time.




FIG. 5.— The flux as a function of view angle ψ for the jet for two case: $\Gamma = 3$ and $\theta_j = 1^\circ$ (thin line); and $\Gamma = 20$ and $\theta_j = 5^\circ$ (thick line).

The second parameter to shape the peaks and dips in the lightcurve is the duty cycle

$$\chi \equiv t_{\text{obs}}/\tau_p < 1, \quad (10)$$

where t_{obs} is the time when the observer's line of sight is inside the jet cone during one precession period. A large value of χ would correspond to narrow dips, while a small χ would correspond to wide dips. This ratio can be obtained by solving the equation $\psi(t) = \theta_j$. In general one obtains $\chi = \chi(\theta_i, \theta_{\text{obs}}, \theta_j)$.

The last parameter

$$\lambda(\theta_i, \theta_{\text{obs}}, \theta_j, \Gamma) \equiv \frac{F(\psi_{\text{max}})}{F(0)} = \frac{F(\theta_i + \theta_{\text{obs}})}{F(0)} < 1, \quad (11)$$

defines how deep the dip is. Besides the three angle parameters $(\theta_i, \theta_{\text{obs}}, \theta_j)$, the Lorentz factor Γ also enters the problem.

Finally, the jet opening angle θ_j and the mean Lorentz factor Γ of Sw J1644+57-like events can be constrained by through a statistical argument. Based on the comparison between observed event rate of these events and the more general TDE event rate constrained from both TDE theory and observational data, one can constrain the beaming factor f_b (Burrows et al. 2011; Lei & Zhang 2011)

$$f_b \sim \max\left(\frac{1}{2\Gamma^2}, \frac{\theta_j^2}{2}\right) \sim \frac{R_{\text{obs}}}{10\%R_{\text{tot}}} \leq 5.5 \times 10^{-3}. \quad (12)$$

This requires that $\Gamma \geq 9.5$ and $\theta_j \leq 6^\circ$. In the following calculations, we take $\theta_j = 6^\circ$ and $\Gamma = 20$ as the typical values. A relative large value of Γ is demanded to reproduce the depth of the observed dips.

The values of θ_i and θ_{obs} can be then obtained by combining Eqs. (10) and (11), if we obtain information about the values of χ and λ . In Fig.6, the relation between θ_i and θ_{obs} for different χ and λ are shown. Based on observations, the value of χ in Sw J1644+57 cannot be too large (e.g., point “d” in Fig.6), since otherwise the dips would be too shallow to be observed (e.g. Fig.7d). Similarly, λ cannot be too small (e.g., point “a” in Fig.6), since otherwise the peaks and dips would be too sharp (e.g. Fig.7a). Observationally, moderate values of both parameters, e.g. $\chi \sim 0.4$ and $\lambda \sim 0.3$, are favored (e.g. points “b” or “c” in Fig.6, see Figs.7b and 7c). Since we observe most of the jet beam, the line of sight cannot be too far away from the BH spin axis, i.e. θ_{obs} is expected to be small. From Fig.6, these constraints suggest that the jet inclination angle θ_i is likely similar to $\sim \theta_j$.

Based on the above discussion, the BP radius R_{BP} could be obtained with the observed jet precession period (2.7 days). Putting all the constrains into the numerical model described in Section 2, we can infer the inclination of the stellar orbit before disruption θ_{orbit} . The results are shown in Fig.8. From Fig.8 we find that the inclination angle of the initial stellar orbit can be as large as 20° . For small BH masses, θ_{orbit} can be much greater than θ_i . However, for large BH masses, θ_{orbit} is not much larger than θ_i , mainly due to a small value of R_T/R_g involved.

Finally, the observed X-ray lightcurve Sw J1644+57 does not have strict periodic dips. In some phases the dips disappear, and in some phases the dips do not appear at exactly the predicted phase (Burrows et al. 2011,




FIG. 6.— The relation between θ_i and θ_{obs} for different χ (thin lines, solid: $\chi = 0.3$; dashed: $\chi = 0.4$; dotted: $\chi = 0.5$; dot-dashed: $\chi = 0.7$) and λ (thick lines, solid: $\lambda = 0.3$; dotted: $\lambda = 0.1$), where the jet opening angle $\theta_j = 6^\circ$ and the Lorentz factor $\Gamma = 20$.

and Fig.1). We believe that this is related to some stochastic processes involved in accretion and jet launching. The Lorentz factor Γ may vary with time, and R_{BP} may slightly vary at different epochs. In order to test this idea, we carry out a range of simulations to the lightcurves. First, we introduce a small time-scale variability in the lightcurve overlapped on the $F(0, t) \propto t^{-5/3}$ envelope (Burrows et al. 2011), which may be a result of stochastic magnetic dissipation (e.g. Zhang & Yan 2011) or fluctuations of accretion rate (e.g. Wang et al. 2006). Next, we introduce a variation of Lorentz factor by allowing Γ to randomly vary in the range of (2, 25) during each precession period. We simulate four cases in Fig.7, which correspond to “a”, “b”, “c” and “d” in Fig.6. We find that the change of Lorentz factor can indeed result in the disappearance of dips at some epochs. The peaks and dips in Fig.7a are well separated and are too spiky. The lightcurve in Fig.7d on the other hand is too smooth, without clear evidence of dips. This suggests that they do not resemble the geometric configuration of Sw J1644+57. The other two cases, i.e., Figs.7b and 7c, give interesting quasi-periodic features with noticeable dips. Even though we cannot fully constrain the geometry of the Sw J1644+57 system, we can say that it should resemble the conditions of these two cases based on the simulation results.

5. CONCLUSIONS AND DISCUSSION

In this paper, we propose that the Sw J1644+57 jet launched from a SMBH precesses. This comes from two independent arguments. First, a tidally disrupted star should not have knowledge about the spin direction of the BH before disruption, so that the original stellar orbit very likely has an inclination with respect to the equatorial plane of the BH. A rapidly spinning BH (which is probably the main reason to launch a relativistic jet, Lei & Zhang 2011) tends to distort the accretion flow and warp the accretion disk due to strong LT frame dragging. This would naturally result in a precessing jet if the accretion rate is low enough. Second, observationally




FIG. 7.— Several simulated XRT lightcurves of Sw J1644+57. The four cases correspond to the “a,b,c,d” points of the parameter space in Fig. 6. (a): $\theta_{\text{obs}} = 5.0$ deg, $\theta_i = 7.4$ deg; (b): $\theta_{\text{obs}} = 4.0$ deg, $\theta_i = 6.0$ deg; (c): $\theta_{\text{obs}} = 2.1$ deg, $\theta_i = 6.3$ deg; (d): $\theta_{\text{obs}} = 3.3$ deg, $\theta_i = 3.3$ deg. It is shown that the case (a) has too sharp dips, while the case (d) essentially smeared out the dips. The cases (b) and (c) would be close to what is observed from the source.




FIG. 8.— The inferred inclination of stellar orbit θ_{orbit} vs. the BH mass, where $\tau_p = 2.7$ days. The solid, dashed and dotted lines correspond to $\theta_i = 7^\circ$, 6° and 5° , respectively.

two pieces of evidence hint towards a precessing jet: the rough 2.7-day periodicity and the noticeable lightcurve dips. We performed a SFC analysis to the observed X-ray lightcurve, and confirmed the 2.7-day rough period. We then carried out detailed theoretical modeling on disk warping and jet precessing, and inferred the underlying parameters of the jet system from the observational data. We found that due to disk warping, the original stellar orbit does not have to be nearly aligned with the BH equatorial plane (cf. Stone & Loeb 2012). Rather, θ_{orbit}

can be as large as 20° . In order to reproduce the depths of the lightcurve dips and their distribution, the Lorentz factor should range from moderate values to values as high as 20. The duty cycle χ , which describes the fraction of time when the line of sight intersects the jet, is found to be a moderate value, slightly smaller than 0.5.

Through simulations, we were able to reproduce lightcurves similar to the observed one. Nixon & King (2012) interprets the diminishing dips in some periods as due to jet nutation. Our SFC analysis does not show another quasi frequency that might be related to nutation. We attribute the diminishing dips to fluctuations in Γ . The mis-match between the 2.7-day period and some dips is also understandable, since the outflow that collimates the jet may not always launched strictly from R_{BP} , or R_{BP} could vary stochastically during the event.

The main uncertainty of our calculation is the viscosity parameters ν_1 and ν_2 . We have assumed a constant value for both throughout the event. More complicated evolution is possible (e.g. Chen et al. 2009). Some efforts to constrain these parameters have been made (e.g. Nixon & King 2012), and future detailed numerical simulations are needed to better understand the viscosity.

We thank other UNLV GRB group members for helpful discussion, and Amanda Maxham for technical assist. This work is supported by NSF under Grant No. AST-0908362, by NASA under Grant No. NNX10AD48G, by National Natural Science Foundation of China (grants 11003004 and 11173011), and National Basic Research Program (“973” Program) of China under Grant No. 2009CB824800. WHL acknowledges a Fellowship from China Scholarship Program for support.

REFERENCES

- Bardeen, J. M., & Petterson, J. A. 1975, *ApJ*, 195, L65
 Bardeen J. M., Press W. H., & Teukolsky S. A. 1972, *ApJ*, 178, 347
 Begelman, M. C., King, A. R., & Pringle, J. E. 2006, *MNRAS*, 370, 399
 Blandford, R. D., & Znajek, R. L. 1977, *MNRAS*, 179, 433
 Blandford, R. D., & Payne, D. G. 1982, *MNRAS*, 199, 882
 Bloom, J. S., et al. 2011, *Science*, 333, 203
 Burrows, D. N., Kennea, J. A., Ghisellini, G., et al. 2011, *Nature*, 476, 421
 Cannizzo, J. K., Troja, E., Lodato, G. 2011, *ApJ*, 742, 32
 Caproni, G., Abraham, Z., Livio, M., & Mosquera Cuesta, H. J. 2007, *MNRAS*, 379, 135
 Chen, L., Wu, S. M., & Yuan, F. 2009, *MNRAS*, 398, 1900
 Fragile, P. C., Blaes, O. M., Anninos, P., & Salmonson, J. D. 2007, *ApJ*, 668, 417
 Gao, H., Lu, Y., & Zhang, S. N. 2010, *ApJ*, 717, 268
 Gao, H., Zhang, B. B., & Zhang, B. 2012, *ApJ*, in press, arXiv:1103.0074
 Gehrels, N. et al. 2004, *ApJ*, 611, 1005
 Giannios, D. & Metzger, B. D. 2012, *MNRAS*, 416, 2102
 Granot, J., Panaitescu, A., Kumar, P., & Woosley, S. E. 2002, *ApJ*, 570, L61
 Greenhill, L. J., Kondratko, P. T., Lovell, J. E. J. et al. 2003, *ApJ*, 582, L11
 Herrnstein, J. R., Greenhill, L. J., & Moran, J. M. 1996, *ApJ*, 468, L17
 Kobayashi, S., Laguna, P., Phinney, E. S. & Mészáros, P. 2004, *ApJ*, 615, 855
 Kinney, A. L., Schmitt, H. R., Clarke, C. J. et al. 2000, *ApJ*, 537, 152
 Lei, W. H., Wang, D. X., Gong, B. P., & Huang, C. Y. 2007, *A&A*, 468, 563
 Lei, W. H., & Zhang, B. 2011, *ApJ*, 740, L27
 Lense, J., & Thirring, H. 1918, *Phys. Z.*, 19, 156
 Levan, A. et al. 2011, *Science*, 333, 199
 Lodato, G., & Pringle, J. E. 2006, *MNRAS*, 368, 1196
 Lodato, G. & Rossi, E. M. 2011, *MNRAS*, 410, 359
 Lu, Y., Huang, Y. F. & Zhang, S. N. 2008, *ApJ*, 684, 1330
 Miyoshi, M., Moran, J., Herrnstein, J., Greenhill, L., Naki, N., Diamond, P., Inoue, M. 1995, *Nature*, 373, 127
 Neufeld, D. A., Maloney, P. R. 1995, *ApJ*, 447, L17
 Nixon, C. J., & King, A. R. 2012, arXiv:1201.1297
 Pringle, J. E. 1982, *MNRAS*, 258, 811
 Scheuer, P. A. G., Feiler, R. 1996, *MNRAS*, 282, 291
 Schmitt, H. R., Pringle, J. E., Clarke, C. J., & Kinney, A. L. 2002, *ApJ*, 575, 150
 Shao, L., Zhang, F. W., Fan, Y. Z., & Wei, D. M. 2011, *ApJ*, 734, L33
 Stone, N., & Loeb, A. 2012, *PRL*, 108, 061302
 Strubbe, L. E. & Quataert, E. 2009, *MNRAS*, 400, 2070
 Wang, D. X., Lei, W. H., & Ye, Y. C. 2006, *ApJ*, 643, 1047
 Wilkins, D. C. 1972, *Phys. Rev. D*, 5, 814
 Zauderer, A. et al. 2011, *Nature*, 476, 425
 Zhang, B. & Yan, H. 2011, *ApJ*, 726, 90
 Zhuravlev, V. V., & Ivanov, P. B. 2008, *MNRAS*, 415, 2122